



**UNIVERSITI PUTRA MALAYSIA**

**CHARACTERIZATION AND GROWTH KINETICS OF LOCAL  
N<sub>2</sub>-FIXING BACTERIUM, *BACILLUS SP.* UPMB10**

**OOI TZE CHEAN**

**FSMB 2002 1**

**CHARACTERIZATION AND GROWTH KINETICS OF LOCAL N<sub>2</sub>-FIXING  
BACTERIUM, *BACILLUS SP.* UPMB10**

**By**

**OOI,TZE CHEAN**

**Thesis Submitted in Fulfillment of the Requirements for the Degree of Master  
of Science in the Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia**

**April 2002**



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

**CHARACTERIZATION AND GROWTH KINETICS OF LOCAL N<sub>2</sub>-FIXING BACTERIUM, *BACILLUS SP.* UPMB10**

**By**

**OOI TZE CHEAN**

**April 2002**

**Chairman : Associate Professor Dr. Arbakariya Ariff**

**Faculty : Food Science and Biotechnology**

The production of locally isolated N<sub>2</sub>-fixing bacteria was undertaken in Fermentation Technology Unit, Enzyme and Microbial Technology Laboratory, Institute of Bioscience, Universiti Putra Malaysia. Cellular studies and biochemical tests conducted on *Bacillus sp* UPMB10 suggests that the bacterium is *Bacillus sphaericus*. Optimization of medium for cultivation of the N<sub>2</sub>-fixing *Bacillus* was achieved using 1.4 g/L of glycerol and 2.0 g/L of yeast extract. Addition of biotin and thiamine did not improve growth of the bacteria. Optimum culture condition for growth of UPMB10 in the 2L: stirred tank fermenter was obtained at initial pH range between pH 6.0-8.0, 30°C, at agitation speed of 600 rpm and airflow rate of 0.5 vvm. Viable cell counts obtained under these conditions were approximately  $3.5 \times 10^9$  cfu/mL.

A model employing the logistic equation was proposed to describe growth of this newly isolated *Bacillus*. The values of the general kinetic parameters were calculated from the analysis of experimental data obtained from a number of culture using batch fermentation. The specific growth rates of  $0.40\text{ h}^{-1}$  and  $0.45\text{ h}^{-1}$  were employed for modeling of bacteria growth in a shake flask and in 2 L fermenter, respectively. The proposed model consisting of general kinetic parameters and the specific growth rate was adequate to describe the fermentation data with sufficient accuracy for prediction of biomass production and substrate consumption.

Due to substrate inhibition, production of the bacteria was further enhanced using an exponential fed-batch fermentation technique. With the specific growth rate maintained at  $0.4\text{ h}^{-1}$ , viable cells obtained using fed-batch fermentation was four times higher than batch cultivation of the bacteria. Cell density and productivity was improved by three fold compared to batch cultivation. In all experiments acetylene reduction assay (ARA) levels remained unchanged and was maintained at  $20\text{ nmol C}_2\text{H}_2/\text{hr/mL}$

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PENCIRIAN DAN KINETIK PERTUMBUHAN BACTERIA PENGIKAT  
NITROGEN TEMPATAN, *BACILLUS SP.* UPMB10**

oleh

**OOI TZE CHEAN**

**April 2002**

**Pengerusi : Professor Madya Dr. Arbakariya Ariff**

**Faculti : Sains Makanan dan Bioteknologi**

Pengeluaran bacteria pengikat nitrogen tempatan telah dijalankan di Unit Teknologi Fermentasi, Makmal Teknologi Enzim dan Microb, Institut Biosains, Universiti Putra Malaysia. Kajian sel dan ujikaji biokimia ke atas bacteria *Bacillus sp* UPMB10 mendapati bahawa bacterium tersebut adalah *Bacillus sphaericus*. Media yang optimum untuk pengkulturan bacilli pengikat nitrogen telah dicapai dengan menggunakan 1.4 g/L glycerol dan 2.0 g/L yeast extract. Penambahan biotina dan thiamina tidak memperbaiki pertumbuhan bacteria. Keadaan fermentasi yang optima telah dicapai dengan pH permulaan di antara pH 6.0-8.0, 30°C dan kadar pemutaran 600 rpm dengan 0.5 vvm pengaliran udara. Anggaran kiraan sel hidup yang diperolehi dalam keadaan demikian adalah  $3.5 \times 10^9$  cfu/mL.

Model menggunakan persamaan logistik telah dicadangkan untuk mewakili pertumbuhan bacilli yang baru ditemui. Nilai parameter kinetik am telah dikira daripada analisis banyak data eksperimen fermentasi sekelompok. Nilai kadar pertumbuhan spesifik  $0.40 \text{ j}^{-1}$  dan  $0.45 \text{ j}^{-1}$  telah digunakan untuk penghasilan model pertumbuhan bakteria dalam 'shake flask' dan dalam fermenter 2 liter. Model dicadangkan yang mengandungi parameter kinetik am dan kadar pertumbuhan spesifik adalah memadai untuk mewakili data fermentasi dengan tepat untuk ramalan penghasilan biojisim dan penggunaan substrat

Oleh kerana perencatan substrat, pengeluaran bakteria boleh ditingkatkan dengan menggunakan teknik fermentasi suapan sekelompok eksponen. Dengan kadar pertumbuhan spesifik ditetapkan pada  $0.4 \text{ j}^{-1}$ , sel hidup yang diperolehi daripada fermentasi suapan sekelompok eksponen adalah empat kali lebih tinggi berbanding pertumbuhan bakteria sekelompok. Kepekatan sel dan produktiviti menggunakan fermentasi suapan sekelompok telah ditingkatkan sebanyak tiga kali berbanding pertumbuhan bakteria sekelompok. Dalam semua eksperimen, kadar penurunan acetylene (ARA) tidak berubah dan tetap pada  $20 \text{ nmol C}_2\text{H}_2/\text{hr/mL}$ .

## ACKNOWLEDGEMENTS

I wish to thank my all postgraduate students and laboratory staff of Fermentation Technology Unit, Enzyme and Microbial Technology Laboratory for their assistance throughout the length of my research. I like to express my deepest gratitude towards my father, mother and also See Pao Theen for always giving me the love and support I need.

I would especially like to thank my supervisor Prof. Madya Dr. Arbakariya b. Ariff and the supervisory committee for their guidance, support and encouragement. Without them I would not have been able to complete my study.

Above all, I praise GOD for His guidance and continuous blessings upon my life

I certify that an Examination Committee met on 2<sup>nd</sup> April 2002 to conduct the final examination of Ooi Tze Chean on his Master of Science thesis entitled "Growth Study and Kinetic Characterization of Local N<sub>2</sub>-Fixing Bacterium *Bacillus* sp. UPMB10" In accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) regulations 1981. The committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are follows:

ISMAIL KARIM, Ph.D.  
Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Chairman)

ARBAKARIYA ARIFF, Ph.D.  
Associate Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)

ZULKIFLI HAJI SHAMSUDDIN, Ph.D.  
Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

HALIMI MOHD. SAUD, Ph.D.  
Lecturer  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

HIRZUN YUSOF, Ph.D.  
Lecturer  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)



SHAMSHEER MOHAMAD RAMADILI, Ph.D.  
Professor  
Deputy Dean, School of Graduate Studies  
Universiti Putra Malaysia

Date: 29 APR 2002



This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirement of the degree of Master of Science

---

AINI IDERIS, Ph.D.  
Professor/Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been dully acknowledge. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

---

**OOI TZE CHEAN**

Date:

## TABLE OF CONTENTS

	Page
ABSTRACT .....	ii
ABSTRAK .....	iv
ACKNOWLEDGMENT .....	vi
APPROVAL SHEETS .....	vii
DECLARATION FORM .....	ix
LISTS OF TABLES .....	xiii
LIST OF FIGURES .....	xv
LIST OF ABBREVIATIONS .....	xix
CHAPTER	
1. INTRODUCTION.....	1
2. LITERATURE REVIEW.....	6
2.1 Nitrogen Cycle.....	6
2.2 Biofertilizer.....	8
2.3 Nitrogen (N <sub>2</sub> )-fixing Bacteria.....	10
2.4 Heterotrophic Free-Living Nitrogen Fixers.....	15
2.5 The Use <i>Bacillus sp.</i> as Biofertilizers.....	17
2.6 The Role of Nitrogenase in N <sub>2</sub> -fixation.....	19
2.7 Acetylene Reduction Assay .....	22
2.8 Carbon Source Requirement of Free-living N <sub>2</sub> Fixers.....	23
2.9 Nitrogen Source Requirement of Free-living N <sub>2</sub> Fixers .....	26
2.10 Carbon-nitrogen Ratio Requirement of Free-living N <sub>2</sub> Fixers .....	28
2.11 Effect of Oxygen on Free-living Nitrogen Fixers .....	28
2.12 Effect of pH on Free-living Nitrogen Fixers....	30
2.13 Effect of Temperature on Free-living Nitrogen Fixers .....	30
2.14 Inoculant Production in Liquid Culture.....	31
2.15 Concluding Remarks.....	34
3. GENERAL MATERIALS AND METHODS.....	35
3.1 Bacterial Strain.....	35
3.2 Inoculum Preparation.....	35
3.3 Medium Composition.....	36
3.4 Experimental Plan.....	37
3.5 Fermenter.....	39
3.6 Analytical Procedures .....	40
3.6.1 Viable Counts .....	40
3.6.2 Measurement of Dry Cell Weight .....	41
3.6.3 Determination of Organic Acids .....	41
3.6.4 Determination of Sugars and Glycerol .....	42
3.6.5 Determination of Total Nitrogen .....	42
3.6.6 Acetylene Reduction Assay .....	42

3.	IDENTIFICATION AND CHARACTERIZATION OF N <sub>2</sub> -FIXING BACTERIUM UPMB10 .....	44
4.1	Introduction .....	44
4.2	Materials and Methods .....	45
4.2.1	Strain and Media .....	45
4.2.2	Cellular Studies .....	45
4.2.3	Cultivation Conditions .....	45
4.2.4	Substrate Utilization .....	46
4.2.5	Additional Analytical Methods .....	46
4.2.6	Cultivation on BATS Medium .....	46
4.2.7	N <sub>2</sub> -fixation and Acetylene Reduction Assay.....	47
4.2.8	Statistical Procedure.....	47
4.3	Result and Discussion .....	47
4.4	Conclusion.....	54
5.	OPTIMIZATION OF MEDIUM FOR BIOMASS PRODUCTION OF N <sub>2</sub> -FIXING <i>BACILLUS SP.</i> UPMB10 .....	56
5.1	Introduction .....	56
5.2	Materials and Methods .....	57
5.2.1	Microorganism and Medium .....	57
5.2.2	Shake Flask Culture .....	57
5.2.3	Analytical Methods .....	58
5.2.4	Statistical Procedure .....	59
5.3	Mathematical Method .....	59
5.4	Results and Discussion .....	61
5.4.1	Effect of Carbon Source .....	61
5.4.2	Effect of Glycerol Concentration .....	65
5.4.3	Effect of Nitrogen Source .....	69
5.4.4	Effect of Yeast Extract Concentration .....	75
5.4.5	Effect of Biotin and Thiamine .....	78
5.5	Conclusion .....	80
6	FERMENTATION CONDITIONS AFFECTING GROWTH OF N <sub>2</sub> -FIXING <i>BACILLUS SP</i> UPMB10 .....	81
6.1	Introduction .....	81
6.2	Materials and Methods .....	82
6.2.1	Microorganism and Medium .....	82
6.2.2	Fermentation in Shake Flasks .....	82
6.2.3	Fermentation in 2L Stirred Tank Bioreactor .....	83
6.2.4	Analytical Methods .....	83
6.3	Results and Discussions .....	86
6.3.1	Effect of Initial pH .....	86
6.3.2	Effect of Temperature .....	89

6.3.3 Effect of Agitation .....	91
6.4 Conclusion .....	96
 7. KINETICS AND MODELLING OF BIOFERTILIZER FERMENTATION USING <i>BACILLUS SP</i> (UPMB10) .....	97
7.1 Introduction .....	97
7.2 Methods .....	99
7.2.1 Microorganism and Medium .....	99
7.2.2 Fermentations .....	99
7.2.3 Analytical Methods .....	100
7.2.4 Development of Mathematical Model .....	100
7.3 Mathematical Methods .....	100
7.3.1 General Balance Equation .....	100
7.3.2 Kinetic Models .....	102
7.3.3 Batch Fermenter Model .....	106
7.4 Results and Discussion .....	108
7.4.1 Batch Fermentation of UPMB10 .....	108
7.4.2 Testing of the Fermentation Models .....	110
7.5 Conclusion .....	112
 8. BATCH AND FED-BATCH CULTIVATION OF <i>BACILLUS SP.</i> (UPMB10) UTILIZING GLYCEROL AS THE CARBON SUBSTRATE FOR GROWTH .....	113
8.1 Introduction .....	113
8.2 Methods .....	114
8.2.1 Microorganism and Inoculum Preparation .....	114
8.2.2 Medium.....	114
8.2.3 Fermentation .....	115
8.2.4 Analytical Methods .....	116
8.3 Theory .....	117
8.3.1 Fed-batch Fermenter Model .....	117
8.4 Result and discussion .....	119
8.4.1 Batch Fermentation .....	119
8.4.2 Fed-batch Fermentation .....	120
8.5 Conclusion .....	126
 9 GENERAL CONCLUSION AND SUGGESTION FOR FURTHER RESEARCH .....	127
 BIBLIOGRAPHY .....	130
 APPENDICES.....	142
 BIOGRAPHICAL SKETCH.....	148

## LISTS OF TABLES

Table		Page
1	Estimates of N <sub>2</sub> fixed by different biofertilizers and technologies availability .....	9
2	The range of N <sub>2</sub> -fixing bacteria .....	11
3	Free-living heterotrophic N <sub>2</sub> -fixing bacteria .....	16
4	Typical carbon sources used for cultivation of free-living diazotrophs .....	26
5	Nitrogen source used for cultivation of different N <sub>2</sub> -fixing bacterial strains .....	27
6	Oxygen requirement for cultivation of different free-living diazotrophs .....	29
7	Culture pH condition effecting growth of free-living N <sub>2</sub> -fixing bacteria .....	30
8	Temperature conditions employed for growth of N <sub>2</sub> -fixing bacteria .....	30
9	Medium for biomass production in <i>Bacillus sphaericus</i> .....	36
10	Bacterial growth of bacterium UPMB10 and some other bacteria on nonselective (NA) and selective (BATS) Media .....	51
11	Comparison between <i>B. sphaericus</i> and bacterium UPMB10.....	55
12	Effect of different carbon sources on growth characteristics of <i>Bacillus sphaericus</i> UPMB10 .....	63
13	Effect of different glycerol concentrations on some growth characteristics of <i>Bacillus sphaericus</i> UPMB10.....	68
14	Effect of different nitrogen sources on some growth characteristics of <i>Bacillus sphaericus</i> UPMB10.....	74
15	Effect of different yeast extract concentrations on some growth characteristics of <i>Bacillus sphaericus</i> .....	78
16	The culture performance of <i>B. sphaericus</i> UPMB10 at different initial pHs .....	88

17	The culture performance of <i>B. sphaericus</i> UPMB10 at different temperatures .....	90
18	The culture performance of <i>B. sphaericus</i> UPMB10 at different agitation speeds .....	92
19	Oxygen uptake rate of UPMB10 under different agitation rates .....	96
20	Comparison of the performance and the kinetic parameter values of UPMB10 during batch culture using the shake flask and the fermenter.....	112
21	Different kinetic parameter values applied during exponential fed-batch fermentation .....	115
22	Growth performance of <i>B. sphaericus</i> UPMB10 in exponential fed-batch fermentation operated at different $\mu$ .....	124
23.	Comparison $\mu$ required for exponential fed-batch fermentation and $\mu$ calculated from experimental data .....	125
24	Growth performance of UPMB10 using batch and fed-batch cultivation techniques .....	126

## LISTS OF FIGURES

Figure		Page
1	The nitrogen cycle .....	7
2	Flowchart of the experimental design .....	38
3	Schematic diagram and dimension of a 2 L stirred tank fermenter.....	39
4	A 2 L stirred tank fermenter .....	40
5	Light microscopy of bacterium UPMB10 .....	48
6	Growth of bacterium UPMB10 on Nutrient Agar (NA).....	49
7	Growth of bacterium UPMB10 in NB at 30°C with different initial pH.....	49
8	Growth of bacterium UPMB10 in NB at initial pH 7.0 under different temperature conditions.....	50
9	Acetylene reduction activity in <i>Bacillus sp.</i> UPMB10, <i>Azospirillum brasiliense Sp7</i> and <i>Bacillus sphaericus</i> 2362.....	53
10a	The profile of viable cell concentration during cultivation of <i>Bacillus sphaericus</i> UPMB10 using different carbon sources.....	62
10b	The profile of residual carbon concentration during the cultivation of <i>Bacillus sphaericus</i> UPMB10 using different carbon sources...	64
10c	The pH profile during cultivation of <i>Bacillus sphaericus</i> UPMB10 using different carbon sources .....	65
11a	Growth profile for cultivation of <i>Bacillus sphaericus</i> UPMB10 using different glycerol concentrations.....	66
11b	Residual carbon concentration obtained during cultivation of <i>Bacillus sphaericus</i> UPMB10 using different glycerol Concentrations.....	67
11c	Culture pH profile during cultivation of <i>Bacillus sphaericus</i> UPMB10 using different glycerol concentrations.....	69



12a	Growth profile for cultivation of <i>Bacillus sphaericus</i> UPMB10 using different amino acids.....	70
12b	Profile of residual nitrogen concentration during cultivation of <i>Bacillus sphaericus</i> UPMB10 using different amino acids .....	70
12c	Culture pH profile during cultivation of UPMB10 using different amino acids .....	71
13a	Profile of viable cell number during cultivation of <i>B. sphaericus</i> UPMB10 using different alternative nitrogen sources.....	73
13b	Profile of residual nitrogen concentration during the cultivation of <i>B. sphaericus</i> UPMB10 using different alternative nitrogen sources.	73
13c	Culture pH profile during cultivation of <i>B. sphaericus</i> UPMB10 using different alternative nitrogen sources .....	74
14a	Profile of viable cell during cultivation of <i>B. sphaericus</i> UPMB10 using different yeast concentrations.....	76
14b	Profile of residual nitrogen concentration during cultivation of <i>B. sphaericus</i> UPMB10 using different yeast extract concentration...	76
14c	Profile of pH during cultivation of <i>B. sphaericus</i> UPMB10 using different yeast extract concentration .....	77
15a	Growth profile during cultivation of <i>B. sphaericus</i> UPMB10 in medium biotin and thiamine .....	79
15b	Culture pH profile during cultivation of <i>B. sphaericus</i> UPMB10 in medium biotin and thiamine.....	79
16	The profile of DOT during dynamic gassing out technique for the measurement of oxygen uptake rate and volumetric oxygen transfer rate ( $K_{La}$ ) .....	85
17a	Growth profile of <i>B. sphaericus</i> UPMB10, during cultivation at different initial pH conditions.....	86
17b	Profile of glycerol concentration during cultivation of <i>B. sphaericus</i> UPMB10 at different initial pHs .....	87
17c	The culture pH profile during cultivation of <i>B. sphaericus</i> UPMB10 at different initial pHs.....	88

18a	Growth profile of <i>B. sphaericus</i> UPMB10 cultivated at different temperatures. ....	89
18b	Profile of glycerol concentration during growth <i>B. sphaericus</i> UPMB10 at different temperatures ....	90
18c	The pH profile during growth of <i>B. sphaericus</i> UPMB10 cultivated at different temperatures ....	91
19a	Growth profile of <i>B. sphaericus</i> UPMB10 cultivated at different agitation speeds ....	93
19b	Profile of glycerol concentration during cultivation of <i>B. sphaericus</i> UPMB10 at different agitation speeds ....	94
19c	Profile of DOT during cultivation of <i>B. sphaericus</i> UPMB10 at different agitation speeds rates.....	94
19d	Profile of pH during cultivation of <i>B. sphaericus</i> UPMB10 at different agitation speeds ....	95
20	Comparison of calculated and experimental data for batch fermentation of <i>B. sphaericus</i> UPMB10 using a shake flask.....	110
21	Comparison of calculated and experimental data for batch fermentation of <i>B. sphaericus</i> UPMB10 in the fermenter.....	111
22	Instrument set-up for fed-batch fermentation ....	116
23	The time course of batch fermentation using glycerol as the carbon source in a 2 L fermenter ....	120
24	The time course of fed-batch fermentation at $\mu = 0.2 \text{ h}^{-1}$ using glycerol as the carbon source in a 2 L fermenter.....	121
25	The time course of fed-batch fermentation at $\mu = 0.25 \text{ h}^{-1}$ using glycerol as the carbon source in a 2 L fermenter.....	121
26	The time course of fed-batch fermentation at $\mu = 0.30 \text{ h}^{-1}$ using glycerol as the carbon source in a 2 L fermenter. ....	122
27	The time course of fed-batch fermentation at $\mu = 0.40 \text{ h}^{-1}$ using glycerol as the carbon source in a 2 L fermenter. ....	122
28	The time course of fed-batch fermentation at $\mu = 0.50 \text{ h}^{-1}$ using glycerol as the carbon source in a 2 L fermenter. ....	123

29	Plot of productivity (P) against specific growth rate ( $\mu$ )	.....	124
30	Plot of logarithmic value of total cell number $\ln[VX]$ against time (t)	.....	125

## LIST OF ABBREVIATIONS

$N_2$	Molecular dinitrogen
C/N	Carbon to nitrogen ratio of medium in mM basis
$D_i$	Impeller diameter
DOT	Dissolved oxygen tension
$\mu_{\max}$	Maximum or initial specific growth rate ( $h^{-1}$ )
$S_i$	Initial substrate concentration (g/L)
$t$	Fermentation time ( $hr^{-1}$ )
$X$	Cell concentration (g/L)
$X_i$	Initial cell concentration (g/L)
$X_m$	Maximum cell concentration (g/L)
$\mu$	Growth rate ( $h^{-1}$ )

## CHAPTER 1

### INTRODUCTION

Although nitrogen gas ( $N_2$ ) makes up 78% of the atmosphere, plants and other microorganisms cannot use it as a source of nitrogen. Nitrogen fixation plays an important role in the conversion of 'inert'  $N_2$  from the atmosphere into usable ammonia. In the twentieth century, increase in crop yield requires that biological  $N_2$  fixation be supplemented increasingly with the use of fixed nitrogen from chemical fertilizers. The development of Haber-Bosch process for catalytically combined atmospheric nitrogen with hydrogen from fossil fuels to produce ammonia has enabled increases in crop yields. This process now accounts for almost all nitrogenous fertilizer production and absorbs approximately 1.5% of the world's energy consumption (Cocking, 2000). In 1990, world consumption of fertilizer nitrogen was about 80 million tons (Brown, 1994).

Nitrogen fertilizer application is essential in Malaysian agriculture as it plays a major role in improving plant growth and yield on highly leached, infertile acid tropical soils. The common sources of nitrogen being used in this country are ammonium sulphate (21% N), ammonium nitrate (26% N), ammonium chloride (25%) and urea (46% N). Vast areas are cultivated with perennial tree crops like oil palm where large quantities of fertilizer are required annually to sustain high crop yields and profitability. At present, Malaysia has a total land area of about 329,733 km<sup>2</sup> while those cultivated with industrial crops, namely, oil palm, rubber, cocoa and paddy, amounted to 5.53 million hectares (5530 km<sup>2</sup>). Of this, oil palm accounts for approximately 50% of the cultivated area

(Raof *et al.*, 1999). In oil palm production alone, the estimated total N fertilizer cost is RM 470 million per year (Amir, 2001).

Recently, environmental and energy concerns has risen from the overuse of nitrogenous fertilizers. Fertilizer nitrogen is normally applied in the zone of diminishing returns in order to maximize yield. There may be as much as 50% inefficiency in terms of plant uptake with resultant nitrate contaminating ground water. Conversion of excess nitrate to nitrous oxide ( $N_2O$ ) by denitrification also produces a greenhouse gas 180 times more potent than carbon dioxide per molecule. It is well known that the Haber-Bosch process is energy expensive. Bockman (1997) pointed out that it takes 1.3 tons of oil to deliver 1 ton of fertilizer nitrogen. At present, the annual cost of fertilizer nitrogen is US\$20-60 billion worldwide (Hardy, 1997). This has highlighted the need for plants to obtain more of their nitrogen from biological  $N_2$  fixation.

Biological  $N_2$  fixation through legume-*Rhizobium* symbiosis sustains soil N balances in temperate regions. In the tropics, however, legumes contribute little to the soil nitrogen economy (Dart, 1986). Non-legumes in association with free-living or associative bacteria types are capable of  $N_2$  fixation. This  $N_2$ -fixing system plays an important role in maintaining N levels in tropical soils. Free-living and associative  $N_2$ -fixing bacteria can be found in the genera *Azospirillum*, *Azotobacter*, *Beijerinckia*, *Derxia*, *Archromobacter*, *Mycobacterium*, *Arthrobacter*, *Bacillus*, *Klebsiella*, *Pseudomonas*, *Clostridium*, *Chromatium*, *Rhodopseudomonas*, *Desulfovibrio*, *Rhodomicrobium*, *Chlorobium* and *Rhodospirillum* (Subba Rao, 1996). Recently, a

newly isolated strain from the genera *Bacillus* (catalogued as UPMB10) has the ability for high rates of N<sub>2</sub>-fixation (Shamsuddin *et al.*, 1999a). This strain has been tested on local oil palms (Amir, 2001) and banana crops (Mia *et al.*, 1999) and was found to significantly enhance their growths. The preliminary results indicate that *Bacillus sp.* (UPMB10) can be used as a potential biological fertilizer to reduce the usage of chemical N fertilizers.

Biofertilizer refers to microorganisms which increase crop growth through biological N<sub>2</sub> fixation, growth-promoting or hormonal substances, increase availability of soil nutrients, and/or control diseases (Ladha, 1997). The application of biofertilizer for improvement of crop yield and productivity has been studied extensively (Dobereiner and Pedrosa, 1987; Okon and Hadar *et al.*, 1987; Arsac, *et al.*, 1990). For the preparation of biofertilizer, large-scale production of starter cultures or microbial inoculants containing either single pure strain or mixed culture is essential before inoculation into suitable solid substrate for composting. In most cases, fermentation deals with a metabolite or a product of the cell. In the case of microbial inoculants, the cell is the product and, accordingly, fermentation should be altered to optimize cell mass production (Bowers, 1982). One of the important approaches for the development of efficient cultivation process for mass production of cell culture is the optimization of medium and cultural conditions. Since starter cultures for biofertilizer are classified as low end products, cheap industrial medium using locally available carbon and nitrogen sources should be used. For aerobic N<sub>2</sub>-fixing bacteria, such as *Azospirillum*, *Azotobacter* and *Bacillus sp.*, it has been found that concentrations of dissolve oxygen (DO) play an important part in determining

biomass yield and N<sub>2</sub>-fixing efficiency of the strain. Optimization may be carried out using systematic experimental design or mathematical models.

Starter culture, containing high-density cells is normally produced by mass cultivation of the microorganism in submerged cultivation using a fermenter. Several modes of fermenter operation, such as batch, continuous and fed-batch cultures are possible for improvement of yield, productivity and reduction of production cost. Although, reports on the application of various modes of fermenter operation for large-scale cultivation of microbial cell are available elsewhere in the literature (e.g. Harwood and Pirt, 1972) this nature of work is scarce for the production of N<sub>2</sub>-fixing microorganisms. Few studies have been devoted to the physiology of N<sub>2</sub>-fixing bacteria in fermenters for biomass production. This is true even for *Azospirillum*, a well-studied N<sub>2</sub> fixer (Fages, 1994). Most of these physiological studies were related with N<sub>2</sub>-fixation activity. For biomass production growth must not be limited by nitrogen, thus a nitrogen source must be added into the medium. Carbon and energy sources, oligoelements and vitamins are also important components of a growth medium.

This study is aimed at the development of an efficient cultivation technique for the production of locally isolated N<sub>2</sub>-fixing *Bacillus sp* (UPMB10) in the fermenter, as microbial inoculants for subsequence use in the preparation of biofertilizers. Thus, the objectives of the study are;

- (i) To conduct basic identification of *Bacillus sp*.



- (ii) To optimize the cultural conditions for enhancement of growth of *Bacillus sp* (UPMB10).
- (iii) To propose models that can be used to describe growth of *Bacillus sp* (UPMB10) and substrate utilization for better understanding of the process.
- (iv) To investigate the feasibility of using a fed-batch fermentation technique for improvement in yield and productivity of the cultivation of *Bacillus sp* (UPMB10) cells.